



Processing Contaminants in Food Production

Granby, Kit; Duedahl-Olesen, Lene; Fromberg, Arvid; Pedreschi, Franco

Publication date:
2011

[Link back to DTU Orbit](#)

Citation (APA):

Granby, K., Duedahl-Olesen, L., Fromberg, A., & Pedreschi, F. (2011). *Processing Contaminants in Food Production*. Abstract from XXV Interamerican Congress of Chemical Engineering, Santiago, Chile.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Processing Contaminants from the Food Production

Kit Granby¹; Lene Duedahl-Olesen¹; Arvid Fromberg¹ and Franco Pedreschi²

¹Technical University of Denmark, The National Food Institute, Food Chemistry, Søborg, Denmark.

²Pontificia Universidad Católica de Chile
Department of Chemical Engineering and Bioprocesses, Santiago

INTRODUCTION

Contaminants like acrylamide, furan or PAHs (polyaromatic hydrocarbons) as e.g. Benz(a)pyrene may be formed during food processing. All of the substances are genotoxic carcinogens, and for that reason mitigation strategies to reduce the levels are needed. Examples of the formation of the processing contaminants and factors that influence the occurrence are given as well as suggestions for mitigation.

EXPERIMENTAL

The analytical methods for detection of acrylamide, furan and PAHs are described in the references 4, 2 and 3 respectively. The analytical techniques performed are LC-MS/MS (acrylamide); headspace GC-MS (furan) and GC-MS (PAHs). The experiments for studying the formation or mitigation of the contaminants are made as lab-scale experiments or in full scale in cooperation with Danish food industries.




Figure 1. Acrylamide (µg/kg) in an extruded breakfast cereal as function of roasting time at usual (Y) or reduced (Y-25°C; Y-50°C).

RESULTS and DISCUSSION

Case 1: acrylamide in breakfast cereal

An industrial case study of a breakfast cereal production showed that the occurrence of acrylamide in the products greatly depends upon the combined effect of temperature and time. The levels increased exponentially with heating time and it is possible to reduce the acrylamide levels by reducing the temperature. By lowering process temperature and prolonging residence time, acrylamide contents were lowered to 20% with the product quality still maintained.




Figure 2. Formation of furan in potato crisps deep-fried at 150-190°C.




Figure 3. Mean and median ΣPAH₂₅ in industrially smoked fish samples

Case 2: Furan in potato crisps

The furan formation of potato crisps produced at similar general conditions as in industry was studied by deep-frying in palm oil at 150°C for 7 min. at 170°C for 5 min or 190°C for 3.5 min. respectively. The different preparation conditions all produce crisps with a water content of approximate 2%. The result from the frying of the home made crisps reveals a linear relationship between the frying temperature and the furan content which at frying conditions 190°C for 3.5 min formed crisps with furan contents up to 50 µg/kg. Hence reducing frying temperature will mitigate furan level in the products.

Case 3: PAHs in smoked fish

The study on ΣPAH₂₅ in industrially smoked fish samples From Danish fishmongers or larger production sites revealed that indirect smoking from a separate smoke generator mitigated the contaminant level compared to direct smoking. The cold smoked products contained less PAHs than the hot smoked product. Furthermore fillets with a larger relative surface contained more PAHs than the whole fish. Increased combustion temperature also increased PAHs.

REFERENCES

- Jensen, BBB.; Lennox, M.; Granby, K. and Adler-Nissen, J. 2008, *Food and Bioprocess Processing* 2008 86 154.
- Fromberg, A.; Fagt, S. and Granby, K. 2009. Furan in heat processed food products including home cooked food products and ready-to-eat products. Scientific Report made for The European Food Safety Authority EFSA-Q-2009-00846, 20 October 2009 (pp. 47)
- Duedahl-Olesen, L.; Christensen, J. H.; Højgård, A.; Granby, K.; Timm-Heinrich, M. *Food Additives & Contaminants A* 2010 27(9) 1294.
- Granby, K.; Nielsen, N.J.; Hedegaard, R.V.; Christensen, T.; Kann, M. and Skibsted, L.H.; 2008. *Food Additives and Contaminants* 25 () 921.